

# Care of the elderly symposium report

EC McGeough

*Specialist Registrar in Medicine for the Elderly, Edinburgh Royal Infirmary, UK*

**ABSTRACT** This symposium covered a wide range of conditions of interest to the geriatrician, the general physician and the general practitioner, including demographic shift, the epidemiology of ageing, diabetes in older people, investigation and management of falls, an update on stroke (including the role of neurovascular clinics, stroke thrombolysis and rehabilitation) and the management of coronary heart disease in the elderly.

**KEYWORDS** Acute coronary syndromes, ageing, clinical guidelines, diabetes mellitus, carotid sinus syndrome, dementia, epidemiology, falls, integrated care, rehabilitation stroke, syncope, transient ischaemic attack, thrombolysis

**DECLARATION OF INTERESTS** No conflict of interests declared.

**Published online July 2009**

**Correspondence to EC McGeough, Medicine for the Elderly Department, Royal Infirmary Edinburgh, Old Dalkeith Road, Edinburgh EH16 4SU, UK**

**tel. +44 (0)131 537 1000**

**e-mail**

**[liz.mcgeough@luht.scot.nhs.uk](mailto:liz.mcgeough@luht.scot.nhs.uk)**

## SESSION I WHO WILL LOOK AFTER ME WHEN I'M OLD?

In his presentation on the epidemiology of ageing, Prof. David Stott (David Cargill Professor of Geriatric Medicine, Royal Infirmary, Glasgow) addressed issues surrounding longevity, the burden of disability and cognitive ageing. The average life expectancy in the developed world is increasing by 0.8 years every decade, and there is also an increase in the number and proportion of older people in our society. It is projected that by 2030, more than a third of the UK population will be over 65 years of age. Healthy lifestyle measures and a higher socio-economic status can lead to a longer life expectancy as well as less disability at the end of life, or a compression of morbidity.<sup>1</sup> Population increases in life expectancy are not linked with any increase in severe disability before death, but are associated with a rise in self-reported chronic disease and moderate/mild disability and self-reported chronic disease. Women live longer, but do seem to have a longer period of disability at the end of life than men.<sup>2</sup> This overall increase in chronic disease burden will lead to higher demand on healthcare service.

Preventable disease, particularly cerebrovascular disease and ischaemic heart disease, are a major contributor to disability and cognitive decline in older age. The Office of Population Censuses and Surveys Library show that more than 30% of over 80-year olds live with severe disability. A third to a half of physical decline in older age is due to a combination of stroke and ischaemic heart disease.<sup>3</sup> Often the underlying pathology is covert, but not benign. The Rotterdam study concluded that 'silent' brain infarction is five times as common as symptomatic infarction in older people.<sup>4</sup> Small vessel disease is a gradual and progressive phenomenon. White matter lesions are associated with slower gait speed, falls in mini mental state examination (MMSE) and slower information processing.<sup>5</sup>

The new concept of cognitive ageing recognises that Alzheimer's dementia and atherosclerosis are

independent pathological processes with common risk factors. It is the combination of these disease processes that lead to cognitive decline and the most significant expression of dementia.<sup>6</sup> We should therefore view cognitive ageing as a continuum and target treatments to actively manage vascular risk and adopt novel approaches in preventing cognitive decline. Diabetes mellitus is an important cause of disability. Dr Terry Aspray (Consultant Geriatrician and Honorary Clinical Senior Lecturer, Sunderland Royal Hospital) outlined the impact of demographic change on the growing epidemic of type 2 diabetes in older age. More than half of the diabetic population is aged over 50 years, and a quarter over 75 years. Moderate glycaemic control in type 2 diabetes leads to longer-term cardiovascular benefit, but there is a paucity of evidence about the optimal diabetic control in older patients as there tended to be to be excluded from clinical trials.

As new drugs for type 2 diabetes are marketed, we are learning more about their effectiveness and side effects in older patients. Rosiglitazone increases fracture risk and other cardiovascular risks<sup>7</sup> and contribute to fluid retention and anaemia. The incretins may be a safe and efficacious option, with confirmed benefits in HbA1c reduction, lower fasting glucose levels and weight loss conferred into the sixth and seventh decades.<sup>8</sup> The gliptins have shown a greater reduction in HbA1c and weight in 70-year olds compared 50-year olds, tend to be safe and well tolerated<sup>9</sup> and can be orally administered.

Complications of diabetes are common in frail older people, particularly in care home residents. Blood pressure- and glucose-lowering strategies reduce macrovascular and microvascular risk. There is strong epidemiological evidence that diabetes is a risk factor for dementia; both hypoglycaemia and hyperglycaemia are linked with impaired cognition. Tight blood pressure control and tight glycaemic control may not always be achievable in the elderly and individualised care may be more appropriate. Dr John Dean (Medical Director for

Quality and Care Improvement and Consultant Diabetologist, NHS Bolton) discussed his model of integrated care for patients with diabetes. Key features of prevention, self care and collaborative care planning between patient, clinician and carer, at significant times of change, with individualised goals and action plans is fundamental. Care planned from a population perspective, with aims to reduce variations in outcomes across practices and geographical areas and lessen current health inequalities, is crucial. This can be achieved by coordinating appropriate roles and extending skills within the primary care, with specialist care teams working across boundaries and providing key support to primary care.

## SESSION 2 FOUND ON THE FLOOR: THE DIAGNOSIS AND MANAGEMENT OF FALLS

Dr Martin Wilson (Consultant Physician, Raigmore Hospital, Inverness) used two scenarios in a highly relevant presentation to demonstrate how the label of 'falls' in the acute receiving unit is a non-specific term. He highlighted the importance of reaching diagnoses in such cases, and illustrated how the all too common mix of acute and chronic pathology can lead to this presenting complaint. Discriminating between acute and chronic issues is essential in understanding the relative loss of function in individuals, and helps avoid common pitfalls in managing this presenting complaint.

In the Stanley Davidson lecture, Prof. Rose Anne Kenny (Head of Department of Medical Gerontology, Trinity College, Dublin) discussed the future of falls management. Differentiating between falls and syncope requires a witnessed account, or an awareness of events. At least 30% of older adults have amnesia for loss of consciousness. Old age, abnormal electrocardiograms, low haematocrit and underlying heart disease are risk factors in predicting hospital admission following acute presentation of falls or syncope.<sup>10</sup> Of all falls, 35% are accidental, 25% have cognitive impairment, 22% have a medical diagnosis (epilepsy or arrhythmia), and 18% are drop attacks.<sup>11</sup> Randomised controlled trials evaluating targeted intervention based on these categories have had positive results.<sup>12</sup> Epilepsy is often considered in the differential of falls or syncope, and is a challenge to diagnose. A series of studies, from all ages, have found that 40% of cases of 'treatment resistant epilepsy' are in fact, due to an underlying cardiovascular abnormality.<sup>13</sup> Increased implantable and wearable cardiac, blood pressure, heart rate and electroencephalogram monitoring, are the future in reaching a diagnosis in falls and syncope.

Guidelines for falls and syncope highlight the importance of screening for falls or gait and balance problems, the indications for multifactorial assessment and individualised multicomponent intervention strategy, and the place of prescribed balance and strengthening programmes.<sup>14</sup>

Carotid sinus syndrome accounts for 30% of syncope and one in four drop attacks. A cardio-inhibitory response in carotid sinus syndrome generally responds to pacemaker insertion. Concomitant lowering of blood pressure associated with the syndrome is present in 30% of cases, and is more challenging to treat. An abnormal carotid sinus response with lowered blood pressure and heart rate is seen with similar prevalence in subsets of adults with and without symptoms.<sup>15</sup> Tau protein deposits found within autonomic cardiovascular nuclei in brainstem in postmortem studies of patients with carotid sinus hypersensitivity may explain why some have a predisposition towards developing the syndrome. Recent studies have suggested a link between cognitive function and the effectiveness of cardiac pacing in carotid sinus syndrome. No reduction in falls or syncope is seen following pacemaker insertion for carotid sinus syndrome in older patients with MMSE <20, comorbid conditions on multiple other medications.<sup>16,17</sup> Longitudinal studies evaluating carotid sinus hypersensitivity as a biomarker for dementia are required, and whether early intervention could prevent falls and cognitive decline.

Falls are common in dementia. From the age of 50 and onwards, memory, concentration, reaction times and executive function have been shown to decline rapidly.<sup>18</sup> The degree of drop in blood pressure, and the time taken for blood pressure to return to baseline on standing, is one of the known predictors of falls in dementia, and a correlation between this blood pressure variability and white matter hyper intensity on magnetic resonance imaging, a marker of cognitive impairment, has been found.<sup>19,20</sup> The causal association between hypotension and secondary neurodegeneration remains unclear. Fundamental research is required to elucidate the underlying mechanisms of falls in cognitively impaired adults so that preventative interventions to be developed.

Finally, Professor Kenny presented the exciting new projects currently under way in her gerontology unit in Dublin, including the Technology Research for Independent Living Research Project (TRIL) and the Irish Life Longitudinal Study on Ageing, which is researching health, wealth and happiness as factors influencing the experience of ageing.

## SESSION 3 I NEED TREATMENT FAST! FUTURE STROKE CARE NOW

Prof. Martin Dennis (Professor of Stroke Medicine, Western General Hospital, Edinburgh) outlined that the main aim of a neurovascular service is to provide secondary prevention and reduce the risk of further stroke. The risk of stroke is highest in the first few days after a transient ischaemic attack (TIA), 10% in the first week on average, and early intervention is necessary to gain the maximum benefit. The ABCD score was developed to stratify absolute risk of stroke after TIA, and is useful in

identifying patients at highest absolute risk of stroke in whom the benefit of treatment will be greatest.<sup>21</sup>

The European model of a daily, accident and emergency department-based TIA service reduces delays to assessment and treatment, but implementation would be a challenge in the UK healthcare system. The EXPRESS (Early use of eXisting PReventative Strategies for Stroke) study found that early intervention significantly reduced subsequent stroke risk.<sup>22</sup> The introduction of the '24/7 TIA hotline' in the Western General Hospital, Edinburgh, in 2007, where telephone consultations occur with an 'on call' stroke specialist, has consequently reduced the average delay to assessment from 11.3 days to 3.3 days. In Scotland, Quality Improvement Scotland standards recommend that 80% of TIAs should be seen within seven days of receipt of referral. Transient ischaemic attack clinics therefore need to be redesigned and remodelled to maximise their benefits and meet these targets. The introduction of modern communication technologies, including telephone hotlines, video conferencing, web-based clinic templates and electronic patient records, all make a major contribution in minimising delays, optimising diagnostic accuracy, improving communication with primary care and providing patients with necessary information.

Prof. Keith Muir (SINAPSE Professor of Clinical Imaging, University of Glasgow) emphasised that the concept of stroke as an evolving process of ischaemic damage occurring over hours and days is now widely recognised. Intervening early to restore blood supply to cortical tissue stops this ischemic process. Intravenous (IV) thrombolysis for ischaemic stroke is an effective treatment that is supported by clinical trials and recommended by national and international guidelines. Thrombolysis with the tissue plasminogen activator alteplase increases the proportion of disability free survivors after stroke, with a number needed to treat of eight or less, when administered within three hours of symptom onset.<sup>23</sup> Recent trial data confirm that IV alteplase is safe and effective up to 4.5 hours after onset of symptoms, but the benefit is time dependent, and declines significantly over the course of these early hours.<sup>24</sup>

Cerebral haemorrhage is a risk of thrombolysis. Clinical trials have used different classification of cerebral haemorrhage following thrombolysis. When tightening the definition of haemorrhage into those that are clinically significant, i.e. large confluent haematoma associated with poorer neurological outcome, there is an incidence of 2.4% or a risk of one in 50 patients treated.<sup>24</sup>

Implementing change requires increased training in recognition of early ischaemic change on computerised tomography scans, a practice that has evolved over the past 15 years. The ASIST study shows that the vast majority of patients do report to hospital within three hours of onset of symptoms, but highlights delays to

appropriate assessment and imaging that frequently occur.<sup>25</sup> By illustrating the service development that has taken place in Glasgow, Prof. Muir demonstrated that services can change substantially, for the better, in response to more focus on hyper-acute stroke care. Rapid recognition, transfer, clinical assessment, imaging and treatment delivery are necessary to optimise clinical outcomes from IV thrombolytic therapy.

Prof. Peter Langhorne (Professor of Stroke Care, Royal Infirmary, Glasgow) discussed stroke rehabilitation. The evidence for organised in-patient (stroke unit) care in improving outcomes in stroke is well established. Rehabilitation is a core characteristic of a stroke unit. Future planning in stroke care will advance the multidisciplinary rehabilitation model to the acute end of stroke care. Systematic review of many observational studies focussing on the implementation of the multi disciplinary stroke unit care model in routine practice has demonstrated that basic models of stroke care across a range of settings, show consistent benefits in more patients surviving, returning home and regaining independence after stroke.<sup>26</sup>

Early supported discharges (ESD) aim to accelerate discharge home from hospital and provide rehabilitation and support in the home setting. Trials have demonstrated a reduction in hospital stay and improved long-term outcome. Services need to be delivered from a well-resourced and coordinated clinical team, and are appropriate for people with mild or moderate stroke.<sup>27</sup>

A substantial number of trials have examined the effectiveness of home-based rehabilitation services provided by therapists, specialist nurses and multidisciplinary teams. Stroke therapy intervention studies show consistent effects in preventing deterioration and maintaining level of activity, although the economic effectiveness of this remains unclear. Similarly, stroke liaison worker interventions seem to improve patient and carer satisfaction, but have not been shown to influence independence or quality of life.

Finally, it is predicted that, by fully implementing and delivering evidence-based stroke services in Scotland, with 10,000 patients suffering a stroke annually, there would be an additional 800 independent survivors per year. Implementing the evidence informed by robust clinical trials, and ensuring optimal service delivery for the majority of stroke patients remains a challenge.

## SESSION 4 IN GOOD HEART – CARDIOLOGY AND THE ELDERLY

Internationally agreed evidence-based guidelines highlight the best care for patients with conditions such as acute coronary syndromes and unstable angina. There are limitations, however, in that the trials that have informed

these guidelines have not included elderly patients. Age is one of the strongest predictors of adverse events, but the elderly are also likely to be at greater risk of the adverse effect of treatment.

Careful evaluation of risks and benefits of individual treatments should be undertaken when determining what is 'best' for an individual patient. The registry-based Grace Risk Score<sup>28</sup> gives prognostic information of mortality or myocardial infarction, following acute coronary syndrome from day zero to hospital discharge and six months. Based on age, heart rate, systolic blood pressure, creatinine, Killip class, ST depression, elevated biomarkers and cardiac arrest, Grace Risk score gives probability percentage of adverse events at these intervals, and may be used to help guide management.

Evidence has accumulated on the efficacy of dual antiplatelet treatment as well as heparin and direct and indirect thrombin inhibitors. Other drugs such as glycoprotein 2b/3a inhibitors also have a role in reducing cardiovascular adverse events. However, the use of multiple agents increases the risk of bleeding, which itself is associated with morbidity and mortality. The CRUSADE registry<sup>29</sup> gives bleeding score and categorises risk of in-hospital major bleed following non-ST elevation myocardial infarction (NSTEMI). Patients on two or more anti thrombotic agents have higher bleeding risk, and a higher risk of major bleed following invasive catheterisation compared to a conservative approach. Bleeding is a strong predictor of mortality, and avoidance of bleeding risk is as important as the desire to reduce thrombotic events.

Dr Mark de Belder (Consultant Cardiologist, James Cook University Hospital, Middlesbrough) outlined the OASIS 5 trial, comparing fondaparinux with enoxaparin.<sup>30</sup> In a complex protocol, this showed no difference in death, myocardial infarction and refractory ischaemia endpoints at day nine, and significant reduction in bleeding with fondaparinux. Fondaparinux showed some mortality benefit in the longer term. The ACUTY trial randomised bivalirudin, a direct antithrombin drug, against heparin and glycoprotein 2b/3a inhibitor, and reported net clinical outcomes in favour of bivalirudin with less associated bleeding.<sup>31</sup> Cardiovascular endpoints of death, myocardial infarction and unplanned revascularisation endpoints were, however, non-significantly in favour of heparin and glycoprotein 2b/3a inhibitors. Thirty-day non-coronary artery bypass graft (CABG) bleeding risk doubles in the over 75-year olds

in the heparin/glycoprotein 2b/3a inhibitor arm when compared with bivalirudin. The TRITON trial randomised patient with ST segment elevation myocardial infarction and NSTEMI to clodidogrel and prasugrel, and found improved efficacy endpoints of death, myocardial infarction, stroke and revascularisation with prasugrel.<sup>32</sup> Prasugrel did, however, have increased associated bleeding. Furthermore, net harm was seen in patients with previous stroke or TIA, and no net benefit was seen in patients over 75 years or with a body weight of less than 60 kg.

Overall, elderly patients should be offered proven treatments, but they need to be informed of the potential risks as well as benefits of treatment. Evaluating when risk outweighs benefits requires clinical judgement.

Dr Andrew Flapan (Consultant Cardiologist, Edinburgh Royal Infirmary) debated the question of coronary intervention. The main purpose of coronary intervention is to improve symptoms and where possible, improve outcome. Data from the Scottish Cardiac Registry from 2003–04 has shown that 10.7% of percutaneous transluminal coronary angioplasty (PTCA) and 12.7% of all CABG procedures were in patients aged over 75 years. Approximately 5% of this patient cohort suffered a major adverse cardiac event following both interventions, percutaneous and surgical intervention, and data at 12 months show a 9.2% and 8.6% mortality following PTCA and CABG respectively. A significant determinant of outcome is whether intervention is delivered as scheduled or unscheduled care. Percutaneous transluminal coronary angioplasty outcomes from the New York registry, in patients aged over 80 years, report a 11.5% death rate and 13.1% major adverse cardiac event rate in unscheduled care, compared with 1.1% death rate and a 1.6% major adverse event rate in scheduled intervention. This is at least a two-fold increase compared with the same data from patients aged 60–80 years.

There is evidence supporting coronary intervention in the elderly population to improve symptoms, but less evidence to support improvement in prognosis. The TIME study<sup>33</sup> found angioplasty to be a useful treatment for stable angina with a significant reduction in hospital admissions and improved quality of life over 12 months, with no mortality benefit. Given the elderly are more likely to suffer complications of coronary intervention, careful consideration should be given prior to undertaking these procedures, with the overall treatment aims established.

## REFERENCES

- 1 Vita AJ, Terry RB, Hubert HB et al. Aging, health risks, and cumulative disability. *N Engl J Med* 1998; 338:1035–41.
- 2 Sauvaget C, Jagger C, Arthur AJ. Active and cognitive impairment-free life expectancies: results from the Melton Mowbray 75+ health checks. *Age Ageing* 2001; 30:509–15.
- 3 Kamper AM, Stott DJ, Hyland M et al; PROSPER Study Group. Predictors of functional decline in elderly people with vascular risk factors or disease. *Age Ageing* 2005; 34:450–5.
- 4 Vermeer SE, Koudstaal PJ, Oudkerk M et al. Prevalence and risk factors of silent brain infarcts in the population-based Rotterdam Scan Study. *Stroke* 2002; 33:21–5.
- 5 Longstreth WT Jr. Brain vascular disease overt and covert. *Stroke* 2005; 36:2062–3.
- 6 Neuropathology Group. Medical Research Council Cognitive Function and Aging Study (MRC CFAS). *Lancet* 2001; 357:169–75.
- 7 Viberti G, Kahn SE, Greene DA et al. A diabetes outcome progression trial (ADOPT): an international multicenter study of the comparative efficacy of rosiglitazone, glyburide, and metformin in recently diagnosed type 2 diabetes. *Diabetes Care* 2002; 25:1737–43.
- 8 Buse JB, Klonoff DC, Nielsen LL et al. Metabolic effects of two years of exenatide treatment on diabetes, obesity, and hepatic biomarkers in patients with type 2 diabetes: an interim analysis of data from the open-label, uncontrolled extension of three double-blind, placebo-controlled trials. *Clin Ther* 2007; 29:139–53.
- 9 Pratley RE, Rosenstock J, Pi-Sunyer FX et al. Management of type 2 diabetes in treatment-naïve elderly patients: benefits and risks of vildagliptin monotherapy. *Diabetes Care* 2007; 30:3017–22.
- 10 Huff JS, Decker WW, Quinn JV et al. Clinical policy: critical issues in the evaluation and management of adult patients presenting to the emergency department with syncope. *Ann Emerg Med* 2007; 49:431–44.
- 11 Davies AJ, Steen N, Kenny RA. Carotid sinus hypersensitivity is common in older patients presenting to an accident and emergency department with unexplained falls. *Age Ageing* 2001; 30:289–93.
- 12 Parry SW, Kenny RA. Drop attacks in older adults: systematic assessment has a high diagnostic yield. *J Am Geriatr Soc* 2005; 53:74–8.
- 13 Zaidi A, Clough P, Cooper P et al. Misdiagnosis of epilepsy: many seizure-like attacks have a cardiovascular cause. *J Am Coll Cardiol* 2000; 36:181–4.
- 14 Guideline for the prevention of falls in older persons. American Geriatrics Society, British Geriatrics Society, and American Academy of Orthopaedic Surgeons Panel on Falls Prevention. *J Am Geriatr Soc* 2001; 49:664–72.
- 15 Kerr SR, Pearce MS, Brayne C et al. Carotid sinus hypersensitivity in asymptomatic older persons: implications for diagnosis of syncope and falls. *Arch Intern Med* 2006; 166:515–20.
- 16 Kenny RA, Seifer C. Brief report – SAFE PACE 2 Syncope And Falls in the Elderly Pacing and Carotid Sinus Evaluation: a randomised control trial of cardiac pacing in older patients with falls and carotid sinus hypersensitivity. *Am J Geriatr Cardiol* 1999; 8:87.
- 17 Parry SW, Steen N, Bexton RS et al. Pacing in elderly recurrent fallers with carotid sinus hypersensitivity (PERF-CSH): a randomised, double-blind, placebo controlled crossover trial. *Heart* 2009; 95:405–9.
- 18 University of Michigan. Health and retirement study: a longitudinal study of health, retirement and aging. Available from: <http://hrsonline.isr.umich.edu>
- 19 Kenny RA, Kalaria R, Ballard C. Neurocardiovascular instability in cognitive impairment and dementia. *Ann NY Acad Sci* 2002; 977:183.
- 20 Kenny RA, Shaw FE, O'Brien JT et al. Carotid sinus syndrome is common in dementia with Lewy bodies and correlates with deep white matter lesions. *J Neurol Neurosurg Psychiatry* 2004; 75:966–71.
- 21 Rothwell PM, Giles MF, Flossman E, et al. A simple score (ABCD) to identify individuals at high early risk of stroke after transient ischaemic attack. *Lancet* 2005; 366:29–36.
- 22 Rothwell PM, Buchan A, Johnston SC. Recent advances in management of transient ischaemic attacks and minor ischaemic strokes. *Lancet Neurol* 2006; 5:323–31.
- 23 Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *N Engl J Med* 1995 333:1581–7.
- 24 Hacke W, Kaste M, Bluhmki E et al. Thrombolysis with alteplase 3 to 4.5 hours after acute ischemic stroke. *N Engl J Med* 2008; 359:1317–29.
- 25 Harraf F, Sharma AK, Brown MM. A multicentre observational study of presentation and early assessment of acute stroke. *BMJ* 2002; 325:17.
- 26 Seenan P, Long M, Langhorne P. Stroke units in their natural habitat. systematic review of observational studies. *Stroke* 2007; 38:1886–92.
- 27 Langhorne P, Holmqvist LVV; Early Supported Discharge Trialists. Early supported discharge after stroke. *Cochrane Database Syst Rev* 2005; (2):CD000443.
- 28 Global Registry of Acute Coronary Events (GRACE) risk score. Available from: <http://www.outcomes-umassmed.org/grace/>
- 29 Subherwal S, Bach RG, Chen AY et al. Baseline risk of major bleeding in non-ST-segment-elevation myocardial infarction: the CRUSADE (Can Rapid risk stratification of Unstable angina patients Suppress ADverse outcomes with Early implementation of the ACC/AHA Guidelines) Bleeding Score. *Circulation* 2009; 119:1873–82.
- 30 Fifth Organization to Assess Strategies in Acute Ischemic Syndromes Investigators, Yusuf S, Mehta SR, Chrolavicius S et al. Comparison of fondaparinux and enoxaparin in acute coronary syndromes. *N Engl J Med* 2006; 354:1464–76.
- 31 Lopes RD, Alexander KP, Manoukian SV et al. Advanced age, antithrombotic strategy, and bleeding in non-ST-segment elevation acute coronary syndromes: results from the ACUITY (Acute Catheterization and Urgent Intervention Triage Strategy) trial. *J Am Coll Cardiol* 2009; 53:1021–30.
- 32 Wiviott SD, Braunwald E, McCabe CH et al. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2007; 357:2001.
- 33 Pfisterer M, Buser P, Osswald S et al. Outcome of elderly patients with chronic symptomatic coronary artery disease with an invasive vs optimized medical treatment strategy: one year results of the randomized TIME trial. *JAMA* 2003; 289:1117–23.